

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SCIENCE

FRIDAY, JULY 19, 1918

CONTENTS	
Transmission of Activation in Passive Metals as a Model of the Protoplasmic or Nervous Type of Transmission: Professor Ralph S.	
LILLIE	51
The Inter-allied Scientific Food Commission	60
Scientific Events:—	
The Asphalt Industry in 1917; Training Camps for Instructors to prepare College Men for Military Service; Guarding Soldiers' Camps against Flies and Mosquitoes; The Weather Bureau and Dr. Cleveland Abbe; Grove Carl Gilbert	62
Scientific Notes and News	65
University and Educational News	67
Discussion and Correspondence:—	
A Musical Cricket-like Chirping of a Grass-hopper: H. A. Allard	67
Scientific Books:—	
Lutz's Field Book of Insects: Professor ALEX. D. MACGILLIVRAY	68
Notes on Meteorology and Climatology:—	
Rainfall in the United States: Dr. Charles F. Brooks	69
Special Articles:—	
A Parallel Mutation in Drosophila Funebris: DR. A. H. STURTEVANT	72
The Kentucky Academy of Science: Alfred	

MSS. intended for publication and books, etc., intended for review should be sent to The Editor of Science, Garrison-on-Hudson, N. Y.

TRANSMISSION OF ACTIVATION IN PASSIVE METALS AS A MODEL OF THE PROTOPLASMIC OR NERV-OUS TYPE OF TRANSMISSION

One of the most remarkable peculiarities of irritable living cells and cellular elements like nerve fibers is the readiness with which chemical or metabolic influence may be transmitted, without accompanying transfer of material, between regions differing in the degree or character of their physiological activity. Thus one region of a muscle or nerve which is in a physiologically more active or "stimulated" state transmits its activity regularly to another more or less distant resting region. The state of activity aroused in the irritable living system by a localized stimulus does not itself remain localized, but tends to spread; the region immediately stimulated imparts a similar state of activity to adjoining regions, these then activate the next adjoining, and in this manner a wave of activation or excitation is propagated over the entire irritable element, often to a long distance from its point of origin. In many cases, as in nerve, there is no decrease in the intensity of the local process as it passes along the element; its characteristics are reduplicated both qualitatively and quantitatively at each point which it reaches in its course; the local excitation is temporary and quickly dies out, each successive region of the tissue becoming active and then returning automatically to its original state of rest. Transmission of this type is known to physiologists as "conduction," and is exhibited in its most highly developed form in the nerves of higher animals. It is, however, by no means peculiar to these structures; any cell or cell element which reacts as a whole to a local stimulus illustrates the same phenomenon; some disturbance affecting the metabolism and functional activity of the living system is radiated from the original